


The transplant is in motion.
Immunosuppressive therapy and rejection risks start moving too.

Viracor gives you the results you need to adjust treatment and keep your patients in the ideal range – minimizing the chance of rejection without compromising the risk for infection.

RISKS:

- Loss of transplant
- GVHD (in BMT)




TEST NAME	TEST CODE	CPT CODE	TURNAROUND TIME
Tacrolimus LC-MS/MS*	30406	80197	Same day (within 12-18 hours from receipt of specimen), Monday through Saturday
Sirolimus LC-MS/MS*	30407	80195	
Everolimus LC-MS/MS*	30409	80169	

* New York Approved

Reduce risks and improve outcomes with faster therapeutic drug level testing.

When timing is critical, can you afford not to know your patient's drug levels?



Relationship between voriconazole dosage and blood levels.

- In this study, up to 35% of patient samples were outside the therapeutic range.
- Detection of voriconazole trough levels outside the therapeutic interval of 1-5.5 mg/L during the first week of therapy may prevent treatment failures and neurological toxicity.

Andres Pascual, Thierry Calandra, Saskia Bolay, Thierry Buclin, Jacques Bille, and Oscar Marchetti, Voriconazole Therapeutic Drug Monitoring in Patients with Invasive Mycoses Improves Efficacy and Safety Outcomes, Clinical Infectious Diseases, 2008;46(2);204 by permission of Infectious Diseases Society of America.

TEST NAME	TEST CODE	CPT CODE	TURNAROUND TIME
Voriconazole LC-MS/MS*	3300	80299	Same day (within 8-12 hours from receipt of specimen), Monday through Saturday
Posaconazole LC-MS/MS*	4200		
Itraconazole LC-MS/MS*	2800		
Isavuconazole LC-MS/MS*	4900		

* New York Approved



Viracor